

Instantons: topological aspects

Marcos Jardim
 IMECC - UNICAMP
 Departamento de Matemática
 Caixa Postal 6065
 13083-970 Campinas-SP, Brazil

February 2, 2008

Abstract

Keywords: Gauge Theory; differential geometry of vector bundles with connections; instantons; anti-self-dual connections; moduli spaces; Yang-Mills equation; Yang-Mills functional; ADHM construction; Nahm's equations; monopoles; Hitchin's equations.

Cross-references: Moduli spaces (62); Instantons in gauge theory (72); Mathematical uses of gauge theory (75); Index theorems (149); Donaldson invariants (255); Characteristic classes (354); Differential geometry (386); Finite dimensional algebras and quivers (418).

1 Introduction

Let X be a closed (connected, compact without boundary) smooth manifold of dimension 4, provided with a Riemannian metric denoted by g . Let Ω_X^p denote space of smooth p -forms on X , i.e. the sections of $\wedge^p TX$. The Hodge operator acting on p -forms:

$$*: \Omega_X^p \rightarrow \Omega_X^{4-p}$$

satisfies $*^2 = (-1)^p$. In particular, $*$ splits Ω_X^2 into two sub-spaces $\Omega_X^{2,\pm}$ with eigenvalues ± 1 :

$$\Omega_X^2 = \Omega_X^{2,+} \oplus \Omega_X^{2,-} . \quad (1)$$

Note also that this decomposition is an orthogonal one, with respect to the inner product:

$$\langle \omega_1, \omega_2 \rangle = \int_X \omega_1 \wedge * \omega_2 .$$

A 2-form ω is said to be self-dual if $*\omega = \omega$ and it is said to be anti-self-dual if $*\omega = -\omega$. Any 2-form ω can be written as the sum

$$\omega = \omega^+ + \omega^-$$

of its self-dual ω^+ and anti-self-dual ω^- components.

Now let E be a complex vector bundle over X as above, provided with a connection ∇ , regarded as a $\mathbb{C}$ -linear operator

$$\nabla : \Gamma(E) \rightarrow \Gamma(E) \otimes \Omega_X^1$$

satisfying the Leibnitz rule:

$$\nabla(f\sigma) = f\nabla\sigma + \sigma \otimes df$$

for all $f \in C^\infty(X)$ and $\sigma \in \Gamma(E)$. Its curvature $F_\nabla = \nabla \circ \nabla$ is a 2-form with values in $\text{End}(E)$, i.e. $F_\nabla \in \Gamma(\text{End}(E)) \otimes \Omega_X^2$, satisfying the Bianchi identity $\nabla F_\nabla = 0$.

The *Yang-Mills equation* is:

$$\nabla * F_\nabla = 0 \tag{2}$$

It is a 2nd-order non-linear equation on the connection ∇ . It amounts to a non-abelian generalization of Maxwell equations, to which it reduces when E is a line bundle; the four components of ∇ are interpreted as the electric and magnetic potentials.

An *instanton* on E is a smooth connection ∇ whose curvature F_∇ is anti-self-dual as a 2-form, i.e. it satisfies:

$$F_\nabla^+ = 0 \quad \text{i.e.} \quad *F_\nabla = -F_\nabla \quad . \tag{3}$$

The instanton equation is still non-linear (it is linear only if E is a line bundle), but it is only 1st-order on the connection.

Note that if F_∇ is either self-dual or anti-self-dual as a 2-form, then the Yang-Mills equation is automatically satisfied:

$$*F_\nabla = \pm F_\nabla \quad \Rightarrow \quad \nabla * F_\nabla = \pm \nabla F_\nabla = 0$$

by the Bianchi identity. In other words, instantons are particular solutions of the Yang-Mills equation. Furthermore, while the Yang-Mills equation (2) makes sense over any Riemannian manifold, the instanton equation (3) is well-defined only in dimension 4.

A gauge transformation is a bundle automorphism $g : E \rightarrow E$ covering the identity. The set of all gauge transformations of a given bundle $E \rightarrow X$ form a group through composition, called the *gauge group* and denoted by $\mathcal{G}(E)$. The gauge group acts on the set of all smooth connections on E by conjugation:

$$g \cdot \nabla = g^{-1} \nabla g \quad .$$

It is then easy to see that (3) is gauge invariant condition, since $F_{g \cdot \nabla} = g^{-1} F_\nabla g$. The anti-self-duality equation (3) is also *conformally invariant*: a conformal change in the metric does not change the decomposition (1), so it preserves self-dual and anti-self-dual 2-forms.

The topological charge k of the instanton ∇ is defined by the integral:

$$\begin{aligned} k &= -\frac{1}{8\pi^2} \int_X \text{tr}(F_\nabla \wedge F_\nabla) \\ &= c_2(E) - \frac{1}{2}c_1(E)^2 \end{aligned} \quad (4)$$

where the second equality follows from Chern-Weil theory.

If X is a smooth, non-compact, complete Riemannian manifold, an instanton on X is an anti-self-dual connection for which the integral (4) converges. Note that in this case, k as above *need not be an integer*; however it is expected to always be quantized, i.e. always a multiple of some fixed (rational) number which depends only on the base manifold X .

Summary. This note is organized as follows. After revisiting the variational approach to the anti-self-duality equation (3), we study instantons over the simplest possible Riemannian 4-manifold, $\mathbb{R}^4$ with the flat Euclidean metric. We present 't Hooft's explicit solutions (Section 3), the ADHM construction (Section 4) and its dimensional reductions to $\mathbb{R}^3$, $\mathbb{R}^2$ and $\mathbb{R}$ (Section 5). We conclude in Section 6 by explaining the construction of the central object of study in gauge theory, the instanton moduli spaces.

2 Variational aspects of Yang-Mills equation.

Given a fixed smooth vector bundle $E \rightarrow X$, let $\mathcal{A}(E)$ be the set of all (smooth) connections on E . The *Yang-Mills functional* is defined by

$$\text{YM} : \mathcal{A}(E) \rightarrow \mathbb{R} \quad (5)$$

$$\text{YM}(\nabla) = \|F_\nabla\|_{L^2}^2 = \int_M \text{tr}(F_\nabla \wedge *F_\nabla)$$

The Euler-Lagrange equation for this functional is exactly the Yang-Mills equation (2). In particular, self-dual and anti-self-dual connections yield critical points of the Yang-Mills functional.

Splitting the curvature into its self-dual and anti-self-dual parts, we have

$$\text{YM}(\nabla) = \|F_\nabla^+\|_{L^2}^2 + \|F_\nabla^-\|_{L^2}^2$$

It is then easy to see that every anti-self-dual connection ∇ is an absolute minimum for the Yang-Mills functional, and that $\text{YM}(\nabla)$ coincides with the topological charge (4) of the instanton ∇ times $8\pi^2$.

One can construct, for various 4-manifolds but most interestingly for $X = S^4$, solutions of the Yang-Mills equations which are neither self-dual nor anti-self-dual. Such solutions do not minimize (5). Indeed, at least for gauge group $SU(2)$ or $SU(3)$, it can be shown that there are no other local minima: any critical point which is neither self-dual nor anti-self-dual is unstable and must be a “saddle point” [3].

3 Instantons on Euclidean space

Let $X = \mathbb{R}^4$ with the flat Euclidean metric, and consider a hermitian vector bundle $E \rightarrow \mathbb{R}^4$. Any connection ∇ on E is of the form $d + A$, where d denotes the usual de Rham operator and $A \in \Gamma(\text{End}(E)) \otimes \Omega_{\mathbb{R}^4}^1$ is a 1-form with values in the endomorphisms of E ; this can be written as follows:

$$A = \sum_{k=1}^4 A_k dx^k \quad , \quad A_k : \mathbb{R}^4 \rightarrow \mathfrak{u}(r) \quad .$$

In the Euclidean coordinates x_1, x_2, x_3, x_4 , the anti-self-duality equation (3) is given by:

$$F_{12} = F_{34} \quad , \quad F_{13} = -F_{24} \quad , \quad F_{14} = F_{23}$$

where

$$F_{ij} = \frac{\partial A_j}{\partial x_i} - \frac{\partial A_i}{\partial x_j} + [A_i, A_j] \quad .$$

The simplest explicit solution is the charge 1 $SU(2)$ -instanton on $\mathbb{R}^4$. The connection 1-form is given by:

$$A_0 = \frac{1}{1 + |x|^2} \cdot \text{Im}(q d\bar{q}) \quad (6)$$

where q is the quaternion $q = x_1 + x_2\mathbf{i} + x_3\mathbf{j} + x_4\mathbf{k}$, while Im denotes the imaginary part of the product quaternion; we are regarding $\mathbf{i}, \mathbf{j}, \mathbf{k}$ as a basis of the Lie algebra $\mathfrak{su}(2)$; from this, one can compute the curvature:

$$F_{A_0} = \left(\frac{1}{1 + |x|^2} \right)^2 \cdot \text{Im}(dq \wedge d\bar{q}) \quad (7)$$

We see that the action density function

$$|F_{A_0}|^2 = \left(\frac{1}{1 + |x|^2} \right)^2$$

has a bell-shaped profile centered at the origin and decaying like r^{-4} .

Let $t_{\lambda, y} : \mathbb{R}^4 \rightarrow \mathbb{R}^4$ be the isometry given by the composition of a translation by $y \in \mathbb{R}^4$ with a homothety by $\lambda \in \mathbb{R}^+$. The pullback connection $t_{\lambda, y}^* A_0$ is still anti-self-dual; more explicitly:

$$A_{\lambda, y} = t_{\lambda, y}^* A_0 = \frac{\lambda^2}{\lambda^2 + |x - y|^2} \cdot \text{Im}(q d\bar{q})$$

$$\text{and } F_{A_{\lambda, y}} = \left(\frac{\lambda^2}{\lambda^2 + |x - y|^2} \right)^2 \cdot \text{Im}(dq \wedge d\bar{q}) \quad .$$

Note that the action density function $|F_A|^2$ has again a bell-shaped profile centered at y and decaying like r^{-4} ; the parameter λ measures the concentration of

the energy density function, and can be interpreted as the “size” of the instanton $A_{\lambda,y}$.

Instantons of topological charge k can be obtained by “superimposing” k basic instantons, via the so-called ‘t Hooft ansatz. Consider the function $\rho : \mathbb{R}^4 \rightarrow \mathbb{R}$ given by:

$$\rho(x) = 1 + \sum_{j=1}^k \frac{\lambda_j^2}{(x - y_j)^2} ,$$

where $\lambda_j \in \mathbb{R}$ and $y_j \in \mathbb{R}^4$. Then the connection 1-form $A = A_\mu dx_\mu$ with coefficients

$$A_\mu = i \sum_{\nu=1}^4 \bar{\sigma}_{\mu\nu} \frac{\partial}{\partial x_\nu} \ln(\rho(x)) \quad (8)$$

is anti-self dual; here, $\bar{\sigma}_{\mu\nu}$ are the matrices given by $(\mu, \nu = 1, 2, 3)$:

$$\bar{\sigma}_{\mu\nu} = \frac{1}{4i} [\sigma_\mu, \sigma_\nu] \quad \bar{\sigma}_{\mu 4} = \frac{1}{2} \sigma_\mu$$

where σ_μ are the Pauli matrices.

The connection (8) correspond to k instantons centered at points y_i with size λ_i . The basic instanton (6) is exactly (modulo gauge transformation) what one obtains from (8) for the case $k = 1$. The ‘t Hooft instantons form a $5k$ parameter family of anti-self-dual connections.

$SU(2)$ -instantons are also the building blocks for instantons with general structure group [2]. Let G be a compact semi-simple Lie group, with Lie algebra $\mathfrak{g}$. Let $\phi : \mathfrak{su}(2) \rightarrow \mathfrak{g}$ be any injective Lie algebra homomorphism. If A is an anti-self-dual $SU(2)$ -connection 1-form, then it is easy to see that $\phi(A)$ is an anti-self-dual G -connection 1-form. Using (8) as an example, we have that:

$$A = i \sum_{\mu,\nu} \phi(\bar{\sigma}_{\mu\nu}) \frac{\partial}{\partial x_\nu} \ln(\rho(x)) dx_\mu \quad (9)$$

is a G -instanton on $\mathbb{R}^4$.

While this guarantees the existence of G -instantons on $\mathbb{R}^4$, note that the instanton (9) might be reducible (e.g. ϕ can simply be the obvious inclusion of $\mathfrak{su}(2)$ into $\mathfrak{su}(n)$ for any n) and that its charge depends on the choice of representation ϕ . Furthermore, it is not clear whether every G -instanton can be obtained in this way, as the inclusion of a $SU(2)$ -instanton through some representation $\phi : \mathfrak{su}(2) \rightarrow \mathfrak{g}$.

4 The ADHM construction

All $SU(r)$ -instantons on $\mathbb{R}^4$ can be obtained through a remarkable construction due to Atiyah, Drinfeld, Hitchin and Manin. It starts by considering hermitian vector spaces V and W of dimension c and r , respectively, and the following data:

$$B_1, B_2 \in \text{End}(V) \quad , \quad i \in \text{Hom}(W, V) \quad , \quad j \in \text{Hom}(V, W) \quad ,$$

so-called ADHM data. Assume moreover that (B_1, B_2, i, j) satisfy the *ADHM equations*:

$$[B_1, B_2] + ij = 0 \quad (10)$$

$$[B_1, B_1^\dagger] + [B_2, B_2^\dagger] + ii^\dagger - j^\dagger j = 0 \quad (11)$$

Now consider the following maps

$$\alpha : V \times \mathbb{R}^4 \rightarrow (V \oplus V \oplus W) \times \mathbb{R}^4$$

$$\beta : (V \oplus V \oplus W) \times \mathbb{R}^4 \rightarrow V \times \mathbb{R}^4$$

given as follows ($\mathbf{1}$ denotes the appropriate identity matrix):

$$\alpha(z_1, z_2) = \begin{pmatrix} B_1 + z_1 \mathbf{1} \\ B_2 + z_2 \mathbf{1} \\ j \end{pmatrix} \quad (12)$$

$$\beta(z_1, z_2) = \begin{pmatrix} -B_2 - z_2 \mathbf{1} & B_1 + z_1 \mathbf{1} & i \end{pmatrix} \quad (13)$$

where $z_1 = x_1 + ix_2$ and $z_2 = x_3 + ix_4$ are complex coordinates on $\mathbb{R}^4$. The maps (12) and (13) should be understood as a family of linear maps parameterized by points in $\mathbb{R}^4$.

A straightforward calculation shows that the ADHM equation (10) imply that $\beta\alpha = 0$ for every $(z_1, z_2) \in \mathbb{R}^4$. Therefore the quotient $E = \ker \beta / \text{im} \alpha = \ker \beta \cap \ker \alpha^\dagger$ forms a complex vector bundle over $\mathbb{R}^4$ of rank r whenever (B_1, B_2, i, j) is such that α is injective and β is surjective for every $(z_1, z_2) \in \mathbb{R}^4$.

To define a connection on E , note that E can be regarded as a sub-bundle of the trivial bundle $(V \oplus V \oplus W) \times \mathbb{R}^4$. So let $\iota : E \rightarrow (V \oplus V \oplus W) \times \mathbb{R}^4$ be the inclusion, and let $P : (V \oplus V \oplus W) \times \mathbb{R}^4 \rightarrow E$ be the orthogonal projection onto E . We can then define a connection ∇ on E through the *projection formula*

$$\nabla s = P \underline{d} \iota(s)$$

where $\underline{d}$ denotes the trivial connection on the trivial bundle $(V \oplus V \oplus W) \times \mathbb{R}^4$. To see that this connection is anti-self-dual, note that projection P can be written as follows

$$P = \mathbf{1} - \mathcal{D}^\dagger \Xi^{-1} \mathcal{D}$$

where

$$\mathcal{D} : (V \oplus V \oplus W) \times \mathbb{R}^4 \rightarrow (V \oplus V) \times \mathbb{R}^4$$

$$\mathcal{D} = \begin{pmatrix} \beta \\ \alpha^\dagger \end{pmatrix}$$

and $\Xi = \mathcal{D} \mathcal{D}^\dagger$. Note that $\mathcal{D}$ is surjective, so that Ξ is indeed invertible. Moreover, it also follows from (11) that $\beta\beta^\dagger = \alpha^\dagger\alpha$, so that $\Xi^{-1} = (\beta\beta^\dagger)^{-1} \mathbf{1}$.

The curvature F_∇ is given by:

$$\begin{aligned} F_\nabla &= P \left(\underline{d}(1 - \mathcal{D}_1^\dagger \Xi^{-1} \mathcal{D}) \underline{d} \right) = P \left(\underline{d} \mathcal{D}^\dagger \Xi^{-1} (\underline{d} \mathcal{D}) \right) = \\ &= P \left((\underline{d} \mathcal{D}^\dagger) \Xi^{-1} (\underline{d} \mathcal{D}) + \mathcal{D}^\dagger \underline{d} (\Xi^{-1} (\underline{d} \mathcal{D})) \right) = (\underline{d} \mathcal{D}^\dagger) \Xi^{-1} (\underline{d} \mathcal{D}) \end{aligned}$$

for $P(\mathcal{D}^\dagger \underline{d}(\Xi^{-1}(\underline{d} \mathcal{D}))) = 0$ on $E = \ker \mathcal{D}$. Since Ξ^{-1} is diagonal, we conclude that F_∇ is proportional to $d\mathcal{D}^\dagger \wedge d\mathcal{D}$, as a 2-form.

It is then a straightforward calculation to show that each entry of $d\mathcal{D}^\dagger \wedge d\mathcal{D}$ belongs to $\Omega^{2,-}$.

The extraordinary accomplishment of Atiyah, Drinfeld, Hitchin and Manin was to show that every instanton, up to gauge equivalence, can be obtained in this way; see e.g. [4]. For instance, the basic $SU(2)$ -instanton (6) is associated with the following data ($c = 1, r = 2$):

$$B_1, B_2 = 0 \quad , \quad i = \begin{pmatrix} 1 \\ 0 \end{pmatrix} \quad , \quad j = \begin{pmatrix} 0 & 1 \end{pmatrix}$$

Remark. The ADHM data (B_1, B_2, i, j) is said to be *stable* if β is surjective for every $(z_1, z_2) \in \mathbb{R}^4$, and it is said to be *costable* if α is injective for every $(z_1, z_2) \in \mathbb{R}^4$. (B_1, B_2, i, j) is *regular* if it is both stable and costable. The quotient:

$$\{\text{regular solutions of (10) and (11)}\} / U(V)$$

coincides with the moduli space of instantons of rank $r = \dim W$ and charge $c = \dim V$ on $\mathbb{R}^4$ (see below). It is also an example of a quiver variety (see the article "Finite dimensional algebras and quivers" by Alistair Savage), associated to the quiver consisting of two vertices V and W , two loop-edges on the vertex V and two edges linking V to W , one in each direction.

5 Dimensional reductions of the anti-self-dual Yang-Mills equation.

As pointed out above, a connection on an hermitian vector bundle $E \rightarrow \mathbb{R}^4$ of rank r can be regarded as 1-form

$$A = \sum_{k=1}^4 A_k(x_1, \dots, x_4) dx^k \quad , \quad A_k : \mathbb{R}^4 \rightarrow \mathfrak{u}(r) \quad .$$

Assuming that the connection components A_k are invariant under translation in one direction, say x_4 , we can think of

$$\underline{A} = \sum_{k=1}^3 A_k(x_1, x_2, x_3) dx^k$$

as a connection on a hermitian vector bundle over $\mathbb{R}^3$, with the fourth component $\phi = A_4$ being regarded as a bundle endomorphism $\phi : E \rightarrow E$, called a Higgs

field. In this way, the anti-self-duality equation (3) reduces to the so-called *Bogomolny (or monopole) equation*:

$$F_{\underline{A}} = *d\phi \quad (14)$$

where $*$ is the Euclidean Hodge star in dimension 3.

Now assume that the connection components A_k are invariant under translation in two directions, say x_3 and x_4 . Consider

$$\underline{A} = \sum_{k=1}^2 A_k(x_1, x_2) dx^k$$

as a connection on a hermitian vector bundle over $\mathbb{R}^2$, with the third and fourth components combined into a complex bundle endomorphism:

$$\Phi = (A_3 + i \cdot A_4)(dx_1 - i \cdot dx_2)$$

taking values on 1-forms. The anti-self-duality equation (3) is then reduced to the so-called *Hitchin's equations*:

$$\begin{cases} F_{\underline{A}} = [\Phi, \Phi^*] \\ \bar{\partial}_{\underline{A}} \Phi = 0 \end{cases} \quad (15)$$

Conformal invariance of the anti-self-duality equation means that Hitchin's equations are well-defined over any Riemann surface.

Finally, assume that the connection components A_k are invariant under translation in three directions, say x_2, x_3 and x_4 . After gauging away the first component A_1 , the anti-self-duality equations (3) reduce to the so-called Nahm's equations:

$$\frac{dT_k}{dx_1} + \frac{1}{2} \sum_{j,l} \epsilon_{kjl} [T_j, T_l] = 0, \quad j, k, l = \{2, 3, 4\} \quad (16)$$

where each T_k is regarded as a map $\mathbb{R} \rightarrow \mathfrak{u}(r)$.

Those interested in monopoles and Nahm's equations are referred to the survey [9] and the references therein. The best source for Hitchin's equations still are Hitchin's original papers [6]. A beautiful duality known as *Nahm transform* relates the various reductions of the anti-self-duality equation to periodic instantons; see the survey article [7].

It is also worth mentioning the book by Mason & Woodhouse [8], where other interesting dimensional reductions of the anti-self-duality equation are discussed, providing a deep relation between instantons and the general theory of integrable systems.

6 The instanton moduli space

Now fix a rank r complex vector bundle E over a 4-dimensional Riemannian manifold X . Observe that the difference between any two connections is a linear

operator:

$$(\nabla - \nabla')(f\sigma) = f\nabla\sigma + \sigma \cdot df - f\nabla'\sigma - \sigma \cdot df = f(\nabla - \nabla')\sigma \quad .$$

In other words, any two connections on E differ by an endomorphism valued 1-form. Therefore, the set of all smooth connections on E , denoted by $\mathcal{A}(E)$, has the structure of an affine space over $\Gamma(\text{End}(E)) \otimes \Omega_M^1$.

The gauge group $\mathcal{G}(E)$ acts on $\mathcal{A}(E)$ via conjugation:

$$g \cdot \nabla := g^{-1} \nabla g \quad .$$

We can form the quotient set $\mathcal{B}(E) = \mathcal{A}(E)/\mathcal{G}(E)$, which is the set of gauge equivalence classes of connections on E .

The set of gauge equivalence classes of anti-self-dual connections on E is a subset of $\mathcal{B}(E)$, and it is called the *moduli space of instantons* on $E \rightarrow X$. The subset of $\mathcal{M}_X(E)$ consisting of irreducible anti-self-dual connections is denoted $\mathcal{M}_X^*(E)$. Since the choice of a particular vector bundle within its topological class is immaterial, these sets are usually labeled by the topological invariants (Chern or Pontrjagyn classes) of the bundle E . For instance, $\mathcal{M}(r, k)$ denotes the moduli space of instantons on a rank r complex vector bundle $E \rightarrow X$ with $c_1(E) = 0$ and $c_2(E) = k > 0$.

It turns out that $\mathcal{M}_X(E)$ can be given the structure of a Hausdorff topological space. In general, $\mathcal{M}_X(E)$ will be singular as a differentiable manifold, but $\mathcal{M}_X^*(E)$ can always be given the structure of a smooth Riemannian manifold. We start by explaining the notion of a L_p^2 vector bundle. Recall that $L_p^2(\mathbb{R}^n)$ denotes the completion of the space of smooth functions $f : \mathbb{R}^n \rightarrow \mathbb{C}$ with respect to the norm:

$$\|f\|_{L_p^2}^2 = \int_X \left(|f|^2 + |df|^2 + \cdots + |d^{(p)}f|^2 \right) \quad .$$

In dimension $n = 4$ and for $p > 2$, by virtue of the Sobolev embedding theorem, L_p^2 consists of continuous functions, i.e. $L_p^2(\mathbb{R}^n) \subset C^0(\mathbb{R}^n)$. So we define the notion of a L_p^2 vector bundle as a topological vector bundle whose transition functions are in L_p^2 , where $p > 2$.

Now fixed a L_p^2 vector bundle E over X , we can consider the metric space $\mathcal{A}_p(E)$ of all connections on E which can be represented locally on an open subset $U \subset X$ as a $L_p^2(U)$ 1-form. In this topology, the subset of irreducible connections $\mathcal{A}_p^*(E)$ becomes an open dense subset of $\mathcal{A}_p(E)$. Since any topological vector bundle admits a compatible smooth structure, we may regard L_p^2 connections as those that differ from a smooth connection by a L_p^2 1-form. In other words, $\mathcal{A}_p(E)$ becomes an affine space modeled over the Hilbert space of L_p^2 1-forms with values in the endomorphisms of E . The curvature of a connection in $\mathcal{A}_p(E)$ then becomes a L_{p-1}^2 2-form with values in the endomorphism bundle $\text{End}(E)$. Moreover, let $\mathcal{G}_{p+1}(E)$ be defined as the topological group of all L_{p+1}^2 bundle automorphisms. By virtue of the Sobolev multiplication theorem, $\mathcal{G}_{p+1}(E)$ has the structure of an infinite dimensional Lie group modeled on a Hilbert space; its Lie algebra is the space of L_{p+1}^2 sections of $\text{End}(E)$.

The Sobolev multiplication theorem is once again invoked to guarantee that the action $\mathcal{G}_{p+1}(E) \times \mathcal{A}_p(E) \rightarrow \mathcal{A}_p(E)$ is a smooth map of Hilbert manifolds. The quotient space $\mathcal{B}_p(E) = \mathcal{A}_p(E)/\mathcal{G}_{p+1}(E)$ inherits a topological structure; it is a metric (hence Hausdorff) topological space. Therefore, the subspace $\mathcal{M}_X(E)$ of $\mathcal{B}_p(E)$ is also a Hausdorff topological space; moreover, one can show that the topology of $\mathcal{M}_X(E)$ does not depend on p .

The quotient space $\mathcal{B}_p(E)$ fails to be a Hilbert manifold because in general the action of $\mathcal{G}_{p+1}(E)$ on $\mathcal{A}_p(E)$ is not free. Indeed, if A is a connection on a rank r complex vector bundle E over a connected base manifold X , which is associated with a principal G -bundle. Then the isotropy group of A within the gauge group:

$$\Gamma_A = \{g \in \mathcal{G}_{p+1}(E) \mid g(A) = A\}$$

is isomorphic to the centralizer of the holonomy group of A within G .

This means that the subspace of irreducible connections $\mathcal{A}_p^*(E)$ can be equivalently defined as the open dense subset of $\mathcal{A}_p(E)$ consisting of those connections whose isotropy group is minimal, that is:

$$\mathcal{A}_p^*(E) = \{A \in \mathcal{A}_p(E) \mid \Gamma_A = \text{center}(G)\} \quad .$$

Now $\mathcal{G}_{p+1}(E)$ acts with constant isotropy on $\mathcal{A}_p^*(E)$, hence the quotient $\mathcal{B}_p^*(E) = \mathcal{A}_p^*(E)/\mathcal{G}_{p+1}(E)$ acquires the structure of a smooth Hilbert manifold.

Remark. The analysis of neighborhoods of points in $\mathcal{B}_p(E) \setminus \mathcal{B}_p^*(E)$ is very relevant for applications of the instanton moduli spaces to differential topology. The simplest situation occurs when A is an $SU(2)$ -connection on a rank 2 complex vector bundle E which reduces to a pair of $U(1)$ and such $[A]$ occurs as an isolated point in $\mathcal{B}_p(E) \setminus \mathcal{B}_p^*(E)$. Then a neighborhood of $[A]$ in $\mathcal{B}_p(E)$ looks like a cone on an infinite dimensional complex projective space.

Alternatively, the instanton moduli space $\mathcal{M}_X(E)$ can also be described by first taking the subset of all anti-self-dual connections and then taking the quotient under the action of the gauge group. More precisely, consider the map:

$$\rho : \mathcal{A}_p(E) \rightarrow L_p^2(\text{End}(E) \otimes \Omega_X^{2,+}) \quad (17)$$

$$\rho(A) = F_A^+$$

Thus $\rho^{-1}(0)$ is exactly the set of all anti-self-dual connections. It is $\mathcal{G}_{p+1}(E)$ -invariant, so we can take the quotient to get:

$$\mathcal{M}_X(E) = \rho^{-1}(0)/\mathcal{G}_{p+1}(E) \quad .$$

It follows that the subspace $\mathcal{M}_X^*(E) = \mathcal{B}_p^*(E) \cap \mathcal{M}_X(E)$ has the structure of a smooth Hilbert manifold. Index theory comes into play to show that $\mathcal{M}_X^*(E)$ is finite-dimensional. Recall that if D is an elliptic operator on a vector bundle over a compact manifold, then D is Fredholm (i.e. $\ker D$ and $\text{coker } D$ are finite dimensional) and its index

$$\text{ind } D = \dim \ker D - \dim \text{coker } D$$

can be computed in terms of topological invariants, as prescribed by the Atiyah-Singer index theorem. The goal here is to identify the tangent space of $\mathcal{M}_X^*(E)$ with the kernel of an elliptic operator.

It is clear that for each $A \in \mathcal{A}_p(E)$, the tangent space $T_A \mathcal{A}_p(E)$ is just $L_p^2(\text{End}(E) \otimes \Omega_X^1)$. We define the pairing:

$$\langle a, b \rangle = \int_X a \wedge *b \quad (18)$$

and it is easy to see that this pairing defines a Riemannian metric (so-called L^2 -metric) on $\mathcal{A}_p(E)$.

The derivative of the map ρ in (17) at the point A is given by:

$$\begin{aligned} d_A^+ : L_p^2(\text{End}(E) \otimes \Omega_X^1) &\rightarrow L_{p-1}^2(\text{End}(E) \otimes \Omega_X^2) \\ a &\mapsto (d_A a)^+ , \end{aligned}$$

so that for each $A \in \rho^{-1}(0)$ we have:

$$T_A \rho^{-1}(0) = \{a \in L_p^2(\text{End}(E)) \otimes \Omega_X^1 \mid d_A^+ a = 0\} .$$

Now for a gauge equivalence class $[A] \in \mathcal{B}_p^*(E)$, the tangent space $T_{[A]} \mathcal{B}_p^*(E)$ consists of those 1-forms which are orthogonal to the fibers of the principal $\mathcal{G}_{p+1}(E)$ bundle $\mathcal{A}_p^*(E) \rightarrow \mathcal{B}_p^*(E)$. At a point $A \in \mathcal{A}_p(E)$, the derivative of the action by some $g \in \mathcal{G}_{p+1}(E)$ is

$$-d_A : L_{p+1}^2(\text{End}(E)) \rightarrow L_p^2(\text{End}(E) \otimes \Omega_X^1) .$$

Usual Hodge decomposition gives us that there is an orthogonal decomposition:

$$L_p^2(\text{End}(E) \otimes \Omega_X^1) = \text{im } d_A \oplus \ker d_A^* ,$$

which means that:

$$T_{[A]} \mathcal{B}_p^*(E) = \{a \in L_p^2(\text{End}(E) \otimes \Omega_X^1) \mid d_A^* a = 0\} .$$

Thus the pairing (18) also defines a Riemannian metric on $\mathcal{B}_p^*(E)$.

Putting these together, we conclude that the space $T_{[A]} \mathcal{M}_X^*(E)$ tangent to $\mathcal{M}_X^*(E)$ at an equivalence class $[A]$ of anti-self-dual connections can be described as follows:

$$T_{[A]} \mathcal{M}_X^*(E) = \{a \in L_p^2(\text{End}(E) \otimes \Omega_X^1) \mid d_A^* a = d_A^+ a = 0\} \quad (19)$$

It turns out that the so-called *deformation operator* $\delta_A = d_A^* \oplus d_A$:

$$\delta_A : L_p^2(\text{End}(E) \otimes \Omega_X^1) \rightarrow L_{p+1}^2(\text{End}(E)) \oplus L_{p-1}^2(\text{End}(E) \otimes \Omega_X^2)$$

is elliptic. Moreover, if A is anti-self-dual then $\text{coker } \delta_A$ is empty, so that $T_{[A]} \mathcal{M}_X^*(E) = \ker \delta_A$. The dimension of the tangent space $T_{[A]} \mathcal{M}_X^*(E)$ is then

simply given by the index of the deformation operator δ_A . Using the Atiyah-Singer index theorem, we have for $SU(r)$ -bundles with $c_2(E) = k$:

$$\dim \mathcal{M}_X^*(E) = 4rk - (r^2 - 1)(1 - b_1(X) + b_+(X)) \quad .$$

The dimension formula for arbitrary gauge group G can be found at [1]. For example, the moduli space of $SU(2)$ instantons on $\mathbb{R}^4$ of charge k is a smooth Riemannian manifold of dimension $8k - 3$. These parameters are interpreted as the $5k$ parameters describing the positions and sizes of k separate instantons, plus $3(k - 1)$ parameters describing their relative $SU(2)$ phases.

The detailed construction of the instanton moduli spaces can be found at [4]. An alternative source is Morgan's lecture notes in [5].

It is interesting to note that $\mathcal{M}_X^*(E)$ inherits many of the geometrical properties of the original manifold X . Most notably, if X is a Kähler manifold, then $\mathcal{M}_X^*(E)$ is also Kähler; if X is a hyperkähler manifold, then $\mathcal{M}_X^*(E)$ is also hyperkähler. One expects that other geometric structures on X can also be transferred to the instanton moduli spaces $\mathcal{M}_X^*(E)$.

References

- [1] M. F. Atiyah, N. J. Hitchin & I. M. Singer. Self-duality in four-dimensional riemannian geometry. *Proc. Royal Soc. London* **362**, 425-461 (1978).
- [2] C. W. Bernard, N. H. Christ, A. H. Guth & E. J. Weinberg. Pseudoparticle parameters for arbitrary gauge groups. *Phys. Rev. D* **16**, 2967-2977 (1977).
- [3] J. P. Bourguignon & H. B. Lawson Jr. Stability and isolation phenomena for Yang-Mills fields *Commun. Math. Phys.* **79** 189-230 (1981).
- [4] S. K. Donaldson & P. B. Kronheimer. *Geometry of four-manifolds*. Oxford: Clarendon Press 1990.
- [5] R. Friedman & J. W. Morgan (eds.) *Gauge theory and the topology of four-manifolds*. Providence, RI: American Mathematical Society 1998.
- [6] N. Hitchin. The self-duality equations on a Riemann surface. *Proc. London Math. Soc.* **55**, 59-126 (1987).
Stable bundles and integrable systems. *Duke Math. J.* **54**, 91-114 (1987).
- [7] M. Jardim. A survey on Nahm transform. *J. Geom. Phys.* **52**, 313-327 (2004).
- [8] L. J. Mason & N. M. J. Woodhouse. *Integrability, self-duality, and twistor theory*. New York, NY: Clarendon Press 1996.
- [9] M. Murray. Monopoles. In: *Geometric analysis and applications to quantum field theory*, 119-135. *Progr. Math.* **205**. Boston, MA: Birkhuser Boston 2002.